

II – Zero-Knowledge Proofs and Applications

David Pointcheval

Ecole normale supérieure/PSL, CNRS & INRIA



Zero-Knowledge Proofs of Knowledge

Introduction

3-Coloring

Examples

Signatures

From Identification to Signature

Forking Lemma

Zero-Knowledge Proofs of Membership

Introduction

Example: DH

Zero-Knowledge Proofs of Knowledge

Zero-Knowledge Proofs of Knowledge

Introduction

3-Coloring

Examples

Signatures


Zero-Knowledge Proofs of Membership

Proof of Knowledge

How do I prove that I know a solution s to a problem P ?

Proof of Knowledge

How do I prove that I know a solution s to a problem P ?



Proof of Knowledge: Soundness


\mathcal{A} knows something. . . What does it mean?

the information can be extracted: **extractor**

Proof of Knowledge: Soundness

\mathcal{A} knows something. . . What does it mean?

the information can be extracted: **extractor**



Proof of Knowledge: Zero-Knowledge

How do I prove that I know a solution s to a problem P ?

I reveal the solution. . .

Proof of Knowledge: Zero-Knowledge

How do I prove that I know a solution s to a problem P ?

I reveal the solution. . .

How can I do it without revealing any information?

Zero-knowledge: **simulation** and **indistinguishability**


Proof of Knowledge: Zero-Knowledge

How do I prove that I know a solution s to a problem P ?

I reveal the solution...

How can I do it without revealing any information?

Zero-knowledge: **simulation** and **indistinguishability**



Zero-Knowledge Proofs of Knowledge

Introduction

3-Coloring


Examples

Signatures

Zero-Knowledge Proofs of Membership


Proof of Knowledge

How do I prove that I know a 3-color covering, without revealing any information?



Proof of Knowledge


How do I prove that I know a 3-color covering, without revealing any information?



I choose a random permutation on the colors

Proof of Knowledge


How do I prove that I know a 3-color covering, without revealing any information?



I choose a random permutation on the colors
and I apply it to the vertices

Proof of Knowledge


How do I prove that I know a 3-color covering, without revealing any information?



I mask the vertices

Proof of Knowledge


How do I prove that I know a 3-color covering, without revealing any information?



I mask the vertices
and send it to the verifier

Proof of Knowledge


How do I prove that I know a 3-color covering, without revealing any information?



The verifier chooses an edge

Proof of Knowledge


How do I prove that I know a 3-color covering, without revealing any information?



The verifier chooses an edge
I open it

Proof of Knowledge

How do I prove that I know a 3-color covering,
without revealing any information?




The verifier chooses an edge

I open it

The verifier checks the validity: 2 different colors


Secure Multiple Proofs of Knowledge: Authentication

If there exists an efficient adversary,
then one can solve the underlying problem:




Secure Multiple Proofs of Knowledge: Authentication

If there exists an efficient adversary,
then one can solve the underlying problem:



Secure Multiple Proofs of Knowledge: Authentication

If there exists an efficient adversary,
then one can solve the underlying problem:



Zero-Knowledge Proofs of Knowledge

Introduction

3-Coloring

Examples

Signatures

Zero-Knowledge Proofs of Membership

3-Pass Zero-Knowledge Proofs

Generic Proof

- Proof of knowledge of x , such that $\mathcal{R}(x, y)$
- \mathcal{P} builds a commitment r and sends it to \mathcal{V}
- \mathcal{V} chooses a challenge $h \xleftarrow{R} \{0, 1\}^k$ for \mathcal{P}
- \mathcal{P} computes and sends the answer s
- \mathcal{V} checks (r, h, s)

Σ -Protocol

- Proof of knowledge of x
- \mathcal{P} sends a commitment r
- \mathcal{V} sends a challenge h
- \mathcal{P} sends the answer s
- \mathcal{V} checks (r, h, s)

3-Pass Zero-Knowledge Proofs

Generic Proof

- Proof of knowledge of x , such that $\mathcal{R}(x, y)$
- \mathcal{P} builds a commitment r and sends it to \mathcal{V}
- \mathcal{V} chooses a challenge $h \xleftarrow{R} \{0, 1\}^k$ for \mathcal{P}
- \mathcal{P} computes and sends the answer s
- \mathcal{V} checks (r, h, s)

Σ -Protocol

- Proof of knowledge of x
- \mathcal{P} sends a commitment r
- \mathcal{V} sends a challenge h
- \mathcal{P} sends the answer s
- \mathcal{V} checks (r, h, s)

Special soundness

If one can answer to two different challenges $h \neq h'$:
 $\implies s$ and s' for a unique r
 \implies one can extract x

- Setting: $n = pq$
 \mathcal{P} knows x , such that $X = x^2 \bmod n$ and wants to prove it to \mathcal{V}
- \mathcal{P} chooses $r \xleftarrow{R} \mathbb{Z}_n^*$, sets and sends $R = r^2 \bmod n$
- \mathcal{V} chooses $b \xleftarrow{R} \{0, 1\}$ and sends it to \mathcal{P}
- \mathcal{P} computes and sends $s = x^b \times r \bmod n$
- \mathcal{V} checks whether $s^2 \stackrel{?}{=} X^b R \bmod n$

One then reiterates t times

For a fixed R , two valid answers s and s' satisfy

$$s^2/X = R = (s')^2 \bmod n \implies X = (s/s')^2 \bmod n$$

And thus $x = s/s' \bmod n \implies$ **Special Soundness**

- Setting: $n = pq$
 \mathcal{P} knows x , such that $X = x^2 \bmod n$ and wants to prove it to \mathcal{V}
- \mathcal{P} chooses $r \xleftarrow{R} \mathbb{Z}_n^*$, sets and sends $R = r^2 \bmod n$
- \mathcal{V} chooses $b \xleftarrow{R} \{0, 1\}$ and sends it to \mathcal{P}
- \mathcal{P} computes and sends $s = x^b \times r \bmod n$
- \mathcal{V} checks whether $s^2 \stackrel{?}{=} X^b R \bmod n$

One then reiterates t times

For a fixed R , two valid answers s and s' satisfy

$$s^2/X = R = (s')^2 \bmod n \implies X = (s/s')^2 \bmod n$$

And thus $x = s/s' \bmod n \implies$ **Special Soundness**

Fiat-Shamir Proof: Extraction

More precisely: the execution of t repetitions depends on


- (b_1, \dots, b_t) from the verifier \mathcal{V}
- ω that (together with the previous b_i ($i < k$)) determines R_k from the prover \mathcal{P}

If $\Pr_{\omega, (b_i)}[\mathcal{V} \text{ accepts } \mathcal{P}] > 1/2^t + \varepsilon$,

there is a good fraction of ω (more than $\varepsilon/2$)

such that $\Pr_{(b_i)}[\mathcal{V} \text{ accepts } \mathcal{S}] \geq 1/2^t + \varepsilon/2$.

For such a good ω : a good node along the successful path



Fiat-Shamir Proof: Simulation

Honest Verifier

Simulation of a triplet: $(R = r^2 \bmod n, b, s = x^b \times r \bmod n)$

for $r \xleftarrow{R} \mathbb{Z}_n^*$ and $b \xleftarrow{R} \{0, 1\}$

Similar to: $(R = s^2/X^b \bmod n, b, s)$

for $s \xleftarrow{R} \mathbb{Z}_n^*$ and $b \xleftarrow{R} \{0, 1\}$

Simulation: random s and b , and set $(R = s^2/X^b \bmod n, b, s)$

Any Verifier

Simulation of a triplet: $(R = r^2 \bmod n, b = \mathcal{V}(\text{view}), s = x^b \times r \bmod n)$

for $r \xleftarrow{R} \mathbb{Z}_n^*$ only!

Similar to: $(R = s^2/X^b \bmod n, b = \mathcal{V}(\text{view}), s)$ for $s \xleftarrow{R} \mathbb{Z}_n^*$

Simulation: random s and β , and set $R = s^2/X^\beta \bmod n$

upon reception of b : if $b = \beta$, output s , else rewind

b and β independent: rewind once over 2 \implies linear time

- Setting: $n = pq$ and an exponent e
 \mathcal{P} knows x , such that $X = x^e \bmod n$ and wants to prove it to \mathcal{V}
- \mathcal{P} chooses $r \xleftarrow{R} \mathbb{Z}_n^*$, sets and sends $R = r^e \bmod n$
- \mathcal{V} chooses $b \xleftarrow{R} \{0, 1\}^t$ and sends it to \mathcal{P}
- \mathcal{P} computes and sends $s = x^b \times r \bmod n$
- \mathcal{V} checks whether $s^e \stackrel{?}{=} X^b R \bmod n$

For a fixed R , two valid answers s and s' satisfy

$$s^e / X^b = R = (s')^e / X^{b'} \bmod n \implies X^{b'-b} = (s'/s)^e \bmod n$$

If e prime and bigger than 2^t , then e and $b' - b$ are relatively prime:

Bezout: $ue + v(b' - b) = 1 \implies X^{v(b'-b)} = (s'/s)^{ve} = X^{1-ue} \bmod n$

As a consequence: $X = ((s'/s)^v X^u)^e \implies$ **Special Soundness**

- Setting: $\mathbb{G} = \langle g \rangle$ of order q
 \mathcal{P} knows x , such that $y = g^{-x}$ and wants to prove it to \mathcal{V}
- \mathcal{P} chooses $k \xleftarrow{R} \mathbb{Z}_q^*$, sets and sends $r = g^k$
- \mathcal{V} chooses $h \xleftarrow{R} \{0, 1\}^t$ and sends it to \mathcal{P}
- \mathcal{P} computes and sends $s = k + xh \bmod q$
- \mathcal{V} checks whether $r \stackrel{?}{=} g^s y^h$

For a fixed r , two valid answers s and s' satisfy

$$g^s y^h = r = g^{s'} y^{h'} \implies y^{h'-h} = g^{s-s'}$$

And thus $x = (s - s')(h' - h)^{-1} \bmod q \implies$ **Special Soundness**

Signatures

Zero-Knowledge Proofs of Knowledge

Signatures

From Identification to Signature

Forking Lemma

Zero-Knowledge Proofs of Membership

Zero-Knowledge Proof

- Proof of knowledge of x
- \mathcal{P} sends a commitment r
- \mathcal{V} sends a challenge h
- \mathcal{P} sends the answer s
- \mathcal{V} checks (r, h, s)

Signature

- Key Generation $\rightarrow (y, x)$
- Signature of $m \rightarrow (r, h, s)$
Commitment r
Challenge $h = \mathcal{H}(m, r)$
Answer s
- Verification of (m, r, s)
compute $h = \mathcal{H}(m, r)$
and check (r, h, s)

Special soundness

If one can answer to two different challenges $h \neq h'$: s and s' for a unique commitment r , one can extract x

Zero-Knowledge Proof

- Proof of knowledge of x
- \mathcal{P} sends a commitment r
- \mathcal{V} sends a challenge h
- \mathcal{P} sends the answer s
- \mathcal{V} checks (r, h, s)

Signature

- Key Generation $\rightarrow (y, x)$
- Signature of $m \rightarrow (r, h, s)$
Commitment r
Challenge $h = \mathcal{H}(m, r)$
Answer s
- Verification of (m, r, s)
compute $h = \mathcal{H}(m, r)$
and check (r, h, s)

Special soundness

If one can answer to two different challenges $h \neq h'$: s and s' for a unique commitment r , one can extract x

Zero-Knowledge Proofs of Knowledge

Signatures

From Identification to Signature

Forking Lemma

Zero-Knowledge Proofs of Membership

The **Forking Lemma** shows an efficient reduction between the signature scheme and the identification scheme, but basically, if an adversary \mathcal{A} produces, with probability $\varepsilon \geq 2/2^k$, a valid signature (m, r, h, s) , then within $T' = 2T$, one gets two valid signatures (m, r, h, s) and (m, r, h', s') , with $h \neq h'$ with probability $\varepsilon' \geq \varepsilon^2/32q_H^3$.

The **special soundness** provides the secret key.

Zero-Knowledge Proofs of Membership

Zero-Knowledge Proofs of Knowledge

Signatures

Zero-Knowledge Proofs of Membership


Introduction

Example: DH

Proof of Membership

How do I prove that a word w lies in a language \mathcal{L} : $P = (w, \mathcal{L})$?

- if $\mathcal{L} \in \mathcal{NP}$: a witness s can help prove that $w \in \mathcal{L}$



If $w \notin \mathcal{L}$:

- Proof (perfect soundness): a powerful \mathcal{A} cannot cheat
- Argument (computational soundness): a limited \mathcal{A} cannot cheat

Proof of Membership

Soundness

$w \in \mathcal{L}$... what does it mean?

a witness **exists**, different from knowing it: no need of extractor

Zero-Knowledge


How do I prove there exists a witness s ? I reveal it...

How can I do it without revealing any information?

Zero-knowledge:

simulation

and **indistinguishability**



Zero-Knowledge Proofs of Knowledge

Signatures

Zero-Knowledge Proofs of Membership

Introduction

Example: DH

Diffie-Hellman Language

In a group $\mathbb{G} = \langle g \rangle$ of prime order q ,

the **DDH**(g, h) assumption states it is hard to distinguish $\mathcal{L} = (u = g^x, v = h^x)$ from $\mathbb{G}^2 = (u = g^x, v = h^y)$

- \mathcal{P} knows x , such that $(u = g^x, v = h^x)$ and wants to prove it to \mathcal{V}
- \mathcal{P} chooses $k \xleftarrow{R} \mathbb{Z}_q^*$, sets and sends $U = g^k$ and $V = h^k$
- \mathcal{V} chooses $h \xleftarrow{R} \{0, 1\}^t$ and sends it to \mathcal{P}
- \mathcal{P} computes and sends $s = k - xh \bmod q$
- \mathcal{V} checks whether $U \stackrel{?}{=} g^s u^h$ and $V \stackrel{?}{=} h^s v^h$

For a fixed (U, V) , two valid answers s and s' satisfy

$$g^s u^h = U = g^{s'} u^{h'} \quad h^s v^h = V = h^{s'} v^{h'}$$

- if one sets $y = (s - s')(h' - h)^{-1} \bmod q \implies u = g^y$ and $v = h^y$
- there exists a witness: **Perfect Soundness**