



Invariant transports of random measures and the extra head problem

Günter Last (Karlsruhe)

joint work with

Peter Mörters (Bath) and Hermann Thorisson (Reykjavik)

presented at the conference

Stochastic Networks and Stochastic Geometry
Paris, January 12-14, 2015

1. Four problems on random shifts

1. Extra head problem

Consider a two-sided sequence of independent and fair coin tosses. Find a coin that landed heads so that the other coin tosses are still independent and fair.

2. Marriage of Lebesgue and Poisson

Let η be a stationary Poisson process in \mathbb{R}^d . Find a point T of η such that

$$\theta_T \eta - \delta_0 \stackrel{d}{=} \eta.$$

3. Poisson matching

Let η and ξ be two independent stationary Poisson processes with equal intensity. Find a point T of ξ such that

$$\theta_T(\eta + \delta_0, \xi) \stackrel{d}{=} (\eta, \xi + \delta_0)$$

4. Unbiased shifts of Brownian motion

Let $B = (B_t)_{t \in \mathbb{R}}$ be a two-sided standard Brownian motion. Find a random time T such that the space-time shifted process $(B_{T+t} - B_T)_{t \in \mathbb{R}}$ is a Brownian motion, independent of B_T .

2. Invariant transports of random measures

Setting

$(\Omega, \mathcal{F}, \mathbb{P})$ is a σ -finite measure space. For the first three problems \mathbb{P} can be taken as probability measure.

Definition

A **random measure** on \mathbb{R}^d is a random element in the space of all locally finite measures on \mathbb{R}^d equipped with the Kolmogorov product σ -field.

Setting

We consider mappings $\theta_s : \Omega \rightarrow \Omega$, $s \in \mathbb{R}^d$, satisfying $\theta_0 = \text{id}_\Omega$ and the **flow** property

$$\theta_s \circ \theta_t = \theta_{s+t}, \quad s, t \in \mathbb{R}^d.$$

The mapping $(\omega, s) \mapsto \theta_s \omega$ is supposed to be measurable. We assume that P is **stationary**, that is

$$P \circ \theta_s = P, \quad s \in \mathbb{R}^d.$$

Definition

A random measure ξ is **invariant** if

$$\xi(\theta_s \omega, B - s) = \xi(\omega, B), \quad \omega \in \Omega, s \in \mathbb{R}^d, B \in \mathcal{B}^d.$$

Definition

Let ξ be an invariant random measure on \mathbb{R}^d . The measure

$$Q_\xi(A) := \iint \mathbf{1}\{\theta_s \omega \in A, s \in B\} \xi(\omega, ds) P(d\omega), \quad A \in \mathcal{F},$$

is called the **Palm measure** of ξ (with respect to P), where $B \in \mathcal{B}^d$ satisfies $0 < \lambda_d(B) < \infty$.

Theorem (Refined Campbell theorem)

Let ξ be an invariant random measure on \mathbb{R}^d . Then

$$E_P \int f(\theta_s, s) \xi(ds) = E_{Q_\xi} \int f(\theta_0, s) ds$$

for all measurable $f : \Omega \times \mathbb{R}^d \rightarrow [0, \infty)$.

Definition

An **allocation rule** is a measurable mapping $\tau : \Omega \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ that is **equivariant** in the sense that

$$\tau(\theta_t \omega, \mathbf{s} - t) = \tau(\omega, \mathbf{s}) - t, \quad \mathbf{s}, t \in \mathbb{R}^d, \text{P-a.e. } \omega \in \Omega.$$

Theorem (L. and Thorisson '09)

Let ξ and η be two invariant random measures with positive and finite intensities. Let τ be an allocation rule and define $T := \tau(\cdot, 0)$. Then

$$Q_\xi(\theta_T \in \cdot) = Q_\eta$$

iff τ is **balancing** ξ and η , that is

$$\int \mathbf{1}\{\tau(\mathbf{s}) \in \cdot\} \xi(d\mathbf{s}) = \eta \quad \text{P-a.e.}$$

Remark

The previous result extends to **weighted transport kernels** and to **LCSC-groups** G ; see L. and Thorisson '09 and L. '10a. It can even be extended to random measures on a space, on which G operates; see L. '10b and Kallenberg '11.

Example

Assume that $\xi = \lambda_d$ is Lebesgue measure and that η is a **simple point process**. An allocation rule τ is balancing ξ and η , iff P-a.e.

$$\lambda_d(C^\tau(t)) = 1, \quad t \in \eta,$$

where the **cell** $C^\tau(t)$ is given by

$$C^\tau(t) := \{\mathbf{s} \in \mathbb{R}^d : \tau(\mathbf{s}) = t\}.$$

Theorem (Holroyd and Peres '05)

Assume that η is a stationary unit-rate Poisson process and let τ be an allocation rule. Then τ is balancing Lebesgue measure and η iff

$$P(\theta_{\tau(0)}\eta \in \cdot) = P(\eta + \delta_0 \in \cdot).$$

Example

Assume that ξ and η are simple point processes. An allocation rule τ is balancing ξ and η , iff τ is a **perfect matching** (P-a.e.) of the points of ξ with the points of η .

Theorem (Holroyd, Pemantle, Peres, Schramm '09)

Assume that ξ and η are independent stationary unit-rate Poisson processes (defined on their canonical probability space) and let τ be an allocation rule. Then τ is balancing ξ and η iff

$$\theta_T(\xi + \delta_0, \eta) \stackrel{d}{=} (\xi, \eta + \delta_0),$$

where $T := \tau((\xi + \delta_0, \eta), 0)$.

3. Local time of Brownian motion

Setting

$B = (B_t)_{t \in \mathbb{R}}$ is a two-sided standard Brownian motion starting in 0 ($B_0 = 0$) defined on its canonical probability space $(\Omega, \mathcal{F}, P_0)$.

Definition

An **unbiased shift** (of B) is a random time T (negative values are allowed) such that:

- $B^{(T)} := (B_{T+t} - B_T)_{t \in \mathbb{R}}$ is a Brownian motion,
- $B^{(T)}$ is independent of B_T .

Example

If $T \geq 0$ is a stopping time, then $(B_{T+t} - B_T)_{t \geq 0}$ is a one-sided Brownian motion independent of B_T . However, the example

$$T := \inf\{t \geq 0: B_t = a\}$$

shows that $(B_{T+t} - B_T)_{t \in \mathbb{R}}$ need not be a two-sided Brownian motion.

Example

Consider a deterministic $T \equiv t_0$. Then $B^{(T)} = (B_{t_0+t} - B_{t_0})_{t \in \mathbb{R}}$ is a two-sided Brownian. However, since $B_{-t_0}^{(T)} = -B_{t_0}$, this two-sided motion is not independent of $B_T = B_{t_0}$.

Remark

An unbiased shift with $B_T = 0$ is characterized by

$$(B_{T+t})_{t \in \mathbb{R}} \stackrel{d}{=} B.$$

According to Mandelbrot ([The Fractal Geometry of Nature](#)) „...the process of Brownian zeros is stationary in a weakened form.“ He is using the (non-rigorous) concept of [conditional stationarity](#).

However, the stopping time

$$T := \inf\{t \geq 1 : B_t = 0\}$$

has the property $B_T = 0$. But clearly $B^{(T)}$ is not a Brownian motion. The missing link will be provided by balancing [local times](#) at different levels.

Definition

Let ℓ^0 be the **local time** (random measure) at zero. Its right-continuous (generalised) inverse is defined as

$$T_r := \begin{cases} \sup\{t \geq 0 : \ell^0[0, t] = r\}, & r \geq 0, \\ \sup\{t < 0 : \ell^0[t, 0] = -r\}, & r < 0. \end{cases}$$

Theorem

Let $r \in \mathbb{R}$. Then T_r is an unbiased shift.

Idea of the proof: The intervals $[T_n, T_{n+1}]$, $n \in \mathbb{Z}$, split B into iid-cycles. The distribution of these cycles is time-reversible.

Definition

The **local time** measure ℓ^x at $x \in \mathbb{R}$ can be defined by

$$\ell^x(C) := \lim_{h \rightarrow 0} \frac{1}{h} \int \mathbf{1}\{s \in C, x \leq B_s \leq x + h\} ds.$$

Hence

$$\int f(B_s, s) ds = \iint f(x, s) \ell^x(ds) dx \quad \mathbb{P}_0\text{-a.s.}$$

for all measurable $f : \mathbb{R}^2 \rightarrow [0, \infty)$.

Definition

For $t \in \mathbb{R}$ the **shift** $\theta_t: \Omega \rightarrow \Omega$ is given by

$$(\theta_t \omega)_s := \omega_{t+s}, \quad s \in \mathbb{R}.$$

For $x \in \mathbb{R}$ let

$$P_x := P_0(B + x \in \cdot), \quad x \in \mathbb{R},$$

where B is the identity on Ω .

Remark

It is possible to choose a **perfect** version of local times, that is a (measurable) kernel satisfying for all $x \in \mathbb{R}$ and P_x -a.e. that ℓ^x is diffuse and

$$\begin{aligned} \ell^x(\theta_t \omega, C - t) &= \ell^x(\omega, C), \quad C \in \mathcal{B}, t \in \mathbb{R}, \\ \ell^x(B, \cdot) &= \ell^0(B - x, \cdot). \end{aligned}$$

Definition

Let

$$P := \int P_x dx$$

be the distribution of a Brownian motion with a „uniformly distributed“ starting value.

Remark

Stationary increments of B imply that P is stationary, that is

$$P = P \circ \theta_s, \quad s \in \mathbb{R}.$$

Theorem (Geman and Horowitz '73)

The Palm (probability) measure of the local time ℓ^x is P_x .

Definition

Let ν be a probability measure on \mathbb{R} . Define

$$P_\nu := \int P_x \nu(dx), \quad \ell^\nu := \int \ell^x \nu(dx).$$

Corollary

P_ν is the Palm probability measure of ℓ^ν .

Remark

In the language of stochastic analysis ℓ^ν is a continuous **additive functional** with **Revuz measure** ν .

4. Existence of unbiased shifts

Definition (Skorokhod embedding problem)

Let ν be a probability measure on \mathbb{R} . A random time T **embeds** ν if B_T has distribution ν .

Theorem

Let T be a random time and ν be a probability measure on \mathbb{R} . Then T is an unbiased shift embedding ν if and only if the allocation rule τ defined by $\tau_T(s) := T \circ \theta_s + s$ is balancing ℓ^0 and ℓ^ν .

Example

Let $r > 0$. Then

$$\tau(s) := \inf\{t > s : \ell^0([s, t]) = r\}, \quad s \in \mathbb{R}.$$

Then τ is an allocation rule balancing ℓ^0 with itself. Hence $T_r = \tau(\cdot, 0)$ is an unbiased shift (embedding δ_0).

Theorem

Let ν be a probability measure on \mathbb{R} with $\nu\{0\} = 0$. Then the stopping time

$$T := \inf\{t > 0: \ell^0[0, t] = \ell^\nu[0, t]\}$$

embeds ν and is an unbiased shift.

Remark

The above stopping time above was introduced in Bertoin and Le Jan (1992) as a solution of the Skorokhod embedding problem.

Theorem (L., Mörters and Thorisson '14)

Let ν be a probability measure on \mathbb{R} . Then there is a non-negative stopping time that is an unbiased shift embedding ν .

Theorem (L., Mörters and Thorisson '14)

Let ξ and η be jointly stationary orthogonal diffuse random measures on \mathbb{R} with finite and equal intensities. Then the mapping $\tau: \Omega \times \mathbb{R} \rightarrow \mathbb{R}$, defined by

$$\tau(s) := \inf\{t > s: \xi[s, t] = \eta[s, t]\}, \quad s \in \mathbb{R},$$

is an allocation rule balancing ξ and η .

Remark

The previous theorem holds in a more general stationary setting. The assumption of equal intensities has to be replaced by

$$\mathbb{E}[\xi[0, 1]|\mathcal{I}] = \mathbb{E}[\eta[0, 1]|\mathcal{I}] \quad \text{P-a.e.},$$

where \mathcal{I} is the **invariant σ -field**. In the Brownian setting, \mathbb{P} is trivial on \mathcal{I} . (If $A \in \mathcal{I}$ then either $\mathbb{P}(A) = 0$ or $\mathbb{P}(A^c) = 0$.)

5. Moment properties of unbiased shifts

Theorem (L., Mörters and Thorisson '14)

If T is an unbiased shift embedding a probability measure $\nu \neq \delta_0$, then

$$E_0 \sqrt{|T|} = \infty.$$

Idea of the proof:

- Take an $x > 0$ such that $\nu[x, \infty) = P(B_T > x) > 0$.
- On the event $\{B_T > x\}$, T can be bounded from below by the minimum of two independent hitting times for $-x$, independent of B_T .
- Use the moment properties of hitting times.

Theorem (L., Mörters and Thorisson '14)

Suppose ν is a distribution with $\nu\{0\} = 0$. If the stopping time $T \geq 0$ is an unbiased shift embedding ν , then

$$E_0 T^{1/4} = \infty.$$

Theorem (L., Mörters and Thorisson '14)

Suppose ν is a distribution with a finite first moment and let T be the Bertoin/Le Jan stopping time. Then, for all $\beta \in [0, 1/4)$,

$$E_0 T^\beta < \infty.$$

Idea of the proof: Recall that

$$T = \inf\{t > 0: X(t) = 0\}$$

where $X_t := \ell^0[0, t] - \ell^\nu[0, t]$. Define a **time-change**

$$U_r := \inf\{t > 0: \ell^0[0, t] + \ell^\nu[0, t] = r\}, \quad r > 0,$$

with respect to a clock which does not tick during the flat pieces of X . Then

$$\tilde{X}(r) := X(U_r), \quad r > 0$$

resembles a random walk whose return times have tails of order $t^{-\frac{1}{2}}$. As $U_r \sim r^2$ by Brownian scaling, the return times for the original X have tails of order $t^{-\frac{1}{4}}$.

6. References

- J. Bertoin and Y. Le Jan (1992). *Ann. Probab.* **20**, 538–548.
- A.E. Holroyd and Y. Peres (2005). Extra heads and invariant allocations. *Ann. Probab.* **33**, 31–52.
- A.E. Holroyd, R. Pemantle, Y. Peres, and O. Schramm (2009). Poisson Matching. *Annales de l'institut Henri Poincaré (B)* **45**, 266–287.
- O. Kallenberg (2011). Invariant Palm and related disintegrations via skew factorization. *Probability Theory and Related Fields* **149**, 279–301.
- G. Last (2010a). Modern random measures: Palm theory and related models. *New Perspectives in Stochastic Geometry*. (W. Kendall und I. Molchanov, eds.). Oxford University Press.

- G. Last (2010b). Stationary random measures on homogeneous spaces. *Journal of Theoretical Probability* **23**, 478–497.
- G. Last, P. Mörters and H. Thorisson (2014). Unbiased shifts of Brownian motion. *Ann. Probab.* **42**, 431–463.
- G. Last and M.P. Penrose (2015). *Lectures on the Poisson Process*. Cambridge University Press, in preparation.
- G. Last and H. Thorisson (2009). Invariant transports of stationary random measures and mass-stationarity. *Ann. Probab.* **37**, 790–813.
- T.M. Liggett (2002). Tagged particle distributions or how to choose a head at random. In *In and Out of Equilibrium* (V. Sidoravicious, ed.) 133–162, Birkhäuser, Boston.
- B. Mandelbrot (1982). *The Fractal Geometry of Nature*. Freeman and Co., San Francisco.
- P. Mörters and I. Redl (2014). Skorokhod embeddings for two-sided Markov chains. arXiv:1407.4734.

Announcement of the Conference

Geometry and Physics of Spatial Random Systems

September 6–11, 2015
Bad Herrenalb (Germany)